



Australian Government



Information on Spikevax (Moderna) COVID-19 vaccine

Last updated: 25 March 2022

About the vaccine

Spikevax (Moderna) is a vaccine that can prevent people from becoming ill from COVID-19. Two doses are required initially (called the primary course). These 2 doses are usually given 4-6 weeks apart. In special circumstances the interval may be longer. The Moderna COVID-19 vaccine can also be used for a booster dose in people aged 18 years and older. The first booster dose is given 3 months or more after the primary course.

Moderna does not contain any live virus, and it cannot give you COVID-19. It contains the genetic code for an important part of the SARS-CoV-2 virus called the spike protein. After getting the vaccine, your body makes copies of the spike protein. Your immune system will then learn to recognise and fight against the SARS-CoV-2 virus, which causes COVID-19. The genetic code is broken down quickly by the body.

Vaccination is voluntary and free. You can discuss any concerns or questions you have about COVID-19 vaccination with your immunisation provider or your GP before you have the vaccine.

Benefits of the vaccine

Very large clinical trials have shown that Moderna is effective in preventing COVID-19 in people aged 12 years and older. Adults who had two doses of Moderna were about 94% less likely to become ill from COVID-19 than people who did not get the vaccine. The vaccine was also effective in people aged over 65 years (86%) and in adolescents aged 12-17 years.

Protection against COVID-19 starts from about 2 weeks after the first dose. While one dose may give some protection, it may only last for the short term. Two doses will give optimal protection. No vaccine is 100% effective, so it is possible that you can still get infected and sick from COVID-19 after vaccination.

The virus that causes COVID-19 could potentially still infect a vaccinated person. Even if they have no symptoms or only mild symptoms, they could still pass it on to others. However, the COVID-19 vaccines currently used in Australia are effective in reducing the likelihood of a vaccinated person transmitting the virus to close contacts if the person is infected.

This is why it is important to continue other preventative measures like:

- physical distancing
- hand washing
- wearing a face mask
- COVID-19 testing and quarantine/isolation as required by your state/territory.

If you have been vaccinated with two doses of Moderna, you should still get a COVID-19 test if you have symptoms that meet testing criteria according to your local health authority (such as fever, cough, sore throat).

Safety of the vaccine

Moderna has been safely given to hundreds of millions of people around the world. The Moderna and Comirnaty (Pfizer) vaccines both have a very rare risk of heart inflammation (myocarditis or pericarditis). This is more commonly seen in males aged under 30 after the second dose. In some countries, myocarditis and pericarditis have been reported more commonly after Moderna than after Pfizer. Most people who have had these conditions after their vaccine have recovered fully. The benefits of vaccination outweigh this very rare risk and vaccination is still recommended for all eligible age groups.

For current information on the frequency and severity of myocarditis and pericarditis following Moderna and Pfizer, please refer to the weekly updates published by the Therapeutic Goods Administration (TGA), available at: www.tga.gov.au/periodic/covid-19-vaccine-weekly-safety-report.

Who can receive this vaccine

People aged 6 years and over can receive Moderna for their primary course.

For more information about vaccination in children aged 6 to 11 years see the Moderna consent form for parents and guardians of children aged 6 to 11 years.

People aged 18 years and older can receive Moderna vaccine for their booster or winter dose

Booster doses

A booster dose refers to an additional vaccine dose after the primary vaccine course. It is intended to strengthen and prolong protection against COVID-19.

People aged 18 and over can have a booster dose of Moderna, 3 months or more after the primary course.

People aged 16 and over can have a booster dose of Pfizer.

Booster doses are not recommended for younger people at this stage.

Some people at high risk of severe illness from COVID-19 are recommended to have an additional winter booster dose from 4 months after the first booster.

For more information on booster and winter doses see: <https://www.health.gov.au/initiatives-and-programs/covid-19-vaccines/getting-your-vaccination/booster-doses>

Staying up to date

To be considered up to date with COVID-19 vaccination, you must have completed all the doses recommended for your age and health status.

Find out about how to stay up to date with COVID-19 vaccines at: <https://www.health.gov.au/initiatives-and-programs/covid-19-vaccines/getting-your-vaccination/stay-up-to-date>

Who should not receive this vaccine

You should not receive this vaccine if you have had:

- **anaphylaxis** (a type of severe allergic reaction) to a previous dose of an mRNA COVID-19 vaccine (i.e. Moderna/Pfizer)
- **anaphylaxis after exposure to any component of the vaccine**, including polyethylene glycol (PEG)
- **any other serious adverse event** that following review by an experienced immunisation provider or medical specialist was attributed to a previous dose of an mRNA COVID-19 vaccine (i.e. Moderna or Pfizer) and without another cause identified.

Precautions for vaccination

People with certain conditions may need additional precautions such as staying for 30 minutes of observation after having their vaccine or consulting an allergy specialist. Tell your immunisation provider if you have had:

- an **allergic reaction to a previous dose** or to an ingredient of an mRNA COVID-19 vaccine (i.e. Moderna or Pfizer)
- **anaphylaxis to other vaccines or to other medicines** – your provider can check to ensure there are no common ingredients with the COVID-19 vaccine you are receiving
- confirmed mastocytosis with recurrent anaphylaxis that requires treatment.

If **you have a bleeding disorder** or you are **taking a blood-thinning medication** (anticoagulant), tell your immunisation provider. Your immunisation provider can help determine whether it is safe for you to have an intramuscular injection and help decide the best timing for injection.

Special circumstances to discuss before vaccination

People with precautionary conditions for Moderna

People with a history of any of the following conditions can receive Moderna but advice should be sought from a GP, immunisation specialist or cardiologist about the best timing of vaccination and whether any additional precautions are recommended:

- recent (i.e. within the past 3 months) or current inflammatory cardiac illness e.g. myocarditis and pericarditis
- acute rheumatic fever or acute rheumatic heart disease (i.e., with active myocardial inflammation)
- acute decompensated heart failure.

Tell your doctor if you had myocarditis or pericarditis diagnosed after a previous dose of Pfizer or Moderna.

People with weakened immune systems (immunocompromise)

People with immunocompromise includes those who have a medical condition that weakens their immune system. It also includes those who may be taking medications that suppress their immune system. Moderna is not a live vaccine. It is safe in people with immunocompromise.

People with severe immunocompromise are recommended to have a third dose of Moderna for their initial course. Severely immunocompromised people aged 5 years and over who received a third primary dose are recommended to receive a booster dose (i.e. 4th dose) at 3 months, in line with the timing of the general population.

People with immunocompromise, including those living with HIV, have a higher risk of severe illness from COVID-19, including a higher risk of death.

Since some people with immunocompromise may have a reduced response to the vaccine, it is important to consider other preventative measures such as physical distancing after vaccination.

Women who are pregnant or breastfeeding

Pregnant women and adolescents should be routinely offered Pfizer or Moderna, at any stage of pregnancy. If you are trying to become pregnant you do not need to delay vaccination or avoid becoming pregnant after vaccination.

Pregnant women with COVID-19 have an increased risk of severe illness and adverse pregnancy outcomes. Real-world evidence has shown that Moderna is safe for pregnant women and breastfeeding women. You can discuss the decision in relation to timing of vaccination with your health professional.

If you are breastfeeding, you can have Moderna. You do not need to stop breastfeeding after vaccination.

Pregnant women aged 16 years or older who received their primary COVID-19 vaccination course 3 or more months ago are recommended to have a booster dose.

People with a history of COVID-19

If you have had COVID-19 in the past, tell your immunisation provider. You can have Moderna after you recover from COVID-19, or may defer for up to 4 months after recovery. If you have ongoing illness from COVID-19, discuss the best timing of vaccination with your treating doctor.

Moderna and children

Moderna has been provisionally approved for use in people aged 6 years or older and cannot be given to younger people.

Ensuring the safety of Moderna

The Therapeutic Goods Administration (TGA) assesses all vaccines in Australia. This ensures that, in order for a vaccine to be approved, it is safe, effective and manufactured to a very high quality standard. A description of the process for approval of COVID-19 vaccines is available at: www.tga.gov.au/covid-19-vaccine-approval-process.

The safety of COVID-19 vaccines will continue to be monitored closely throughout the COVID-19 vaccination program.

You can report suspected side effects to your vaccination provider or other healthcare professional. They will then make a formal report on your behalf to your state or territory health department or directly to the TGA.

If you would prefer to report it yourself, please visit www.tga.gov.au/reporting-suspected-side-effects-associated-covid-19-vaccine and follow the directions on the page.